



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

EP99/8323

REC'D 23 DEC 1999

WIPO PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

09/807172

Patentanmeldung Nr. Patent application No. Demande de brevet n°

99303118.6

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN
THE HAGUE, 17/12/99
LA HAYE, LE



ANTIGEN-BINDING PROTEINSFIELD OF THE INVENTION

5 The present invention relates to the preparation of multivalent and multispecific binding proteins. In particular, the invention relates to the preparation of antigen binding proteins comprising a plurality of binding units linked in series by means of intervening polypeptide linker groups, the amino acid sequence of which linker groups confer
10 restricted conformational flexibility.

BACKGROUND OF THE INVENTION

15 There is considerable interest in the preparation of multivalent and/or multispecific antigen binding proteins. Antigen binding proteins which are multivalent (that is, comprise more than one antigen binding site), more especially those which are also multispecific (where the antigen binding sites have differing antigen specificities) have found particular application in the fields of
20 diagnosis or therapy, for example, where the construction of binding proteins having binding activity against both target site and diagnostic or therapeutic agent allows for targeted delivery of the diagnostic or therapeutic agent to the intended site of action. Other uses for which multivalent and multispecific binding proteins have
25 been proposed include assays, such as immunoassays and agglutination assays, and purification processes.

Those multivalent, multispecific antigen binding proteins which have been described in the literature to date rely, in general, on the
30 association of antibody light and heavy chain variable domains for the formation of the antigen binding site.

Thus, constructs comprising two or more polypeptide chains are described in WO 94/09131 (Scotgen Limited) and WO 97/14719 (Unilever)
35 and WO 97/38102 (Unilever); multivalent molecules comprising two or more single chain Fv molecules linked together are described in WO 93/11161 (Enzon Inc.) and WO 94/13806 (Dow Chemical Co.).

WO 94/04678 (Casterman et al) describes immunoglobulins capable of

exhibiting the functional properties of classical, four chain, immunoglobulins but which comprise two heavy polypeptide chains only and are naturally devoid of light polypeptide chains. Fragments corresponding to isolated VH domains or to VH dimers linked by the hinge disulphide are also disclosed. These immunoglobulins, which may be isolated from Camelids, do not rely on the association of heavy and light chain variable domains for the formation of the antigen-binding site; instead, the heavy chain variable domain (hereinafter VHH) alone forms the complete antigen binding site, constituting a single domain binding site.

In their later patent application, WO 96/34103, Casterman et al disclose multivalent, multispecific constructs comprising VHH fragments combined with a linker sequence. Suitable linker sequences disclosed and exemplified are derived from sequences corresponding to the hinge domain of an immunoglobulin devoid of light chains.

In the Applicant's co-pending patent application number PCT/EP98/06991, filed 26th October 1998, there are disclosed multivalent, multispecific antigen-binding proteins comprising a polypeptide comprising in series two or more single domain binding units which are preferably variable domains of a heavy chain derived from an immunoglobulin naturally devoid of light chains. The individual single domain binding units may suitably be linked by means of peptide linkers, preferably flexible peptide linkers, which allow the variable domains to flex in relation to each other with the aim of ensuring that they can bind to multiple antigenic determinants simultaneously.

There remains a continuing need for the development of improved methods for producing multivalent and/or multispecific binding proteins, especially antigen binding proteins. In particular, there is commercial interest in producing molecules which not only have improved binding activity but which also can be produced economically on a large scale.

SUMMARY OF THE INVENTION

In a first aspect, the invention provides the use of a polypeptide

group, the amino acid sequence of which group confers restricted conformational flexibility, as a linking group to link binding units in a multivalent binding protein.

- 5 The invention also provides a multivalent binding protein comprising a plurality of binding units linked by means of intervening polypeptide linker groups, the amino acid sequence of which linker group confers ~~restricted conformational flexibility.~~

- 10 The invention further provides a nucleotide sequence encoding a multivalent antigen binding protein according to the invention and cloning and expression vectors comprising such nucleotide sequences. Also provided are host cells transformed with vectors comprising such nucleotide sequences.

15

As used herein, a 'multivalent binding protein' is a protein which has more than one binding units which allow for specific binding with a molecule partner in a binding pair. Included within this are bivalent, trivalent and so on. Examples of suitable binding units
20 include antigen binding domains of antibodies, binding domains of receptors such as hormone receptors, lectins, enzymes, and cell adhesion molecules. A 'multivalent antigen binding protein' is a protein which has more than one antigen binding unit.

- 25 An 'antigen binding unit' is any structure which exhibits antigen-binding activity. This may be an antibody or an immunologically active fragment thereof. An 'antibody' refers to an immunoglobulin which may be derived from natural sources or synthetically produced. Unless indicated otherwise, 'antibody' and 'immunoglobulin' are used
30 synonymously throughout this specification.

An antibody fragment is a portion of a whole antibody which retains the ability to exhibit antigen-binding activity. The antigen binding site may be formed through association of antibody light and heavy
35 chain variable domains or may comprise individual antibody variable domains, constituting a single domain binding site.

Suitable fragments include Fab (comprising an antibody light chain associated with the V_H and C_{H1} domains of an antibody heavy chain), Fv

(comprising the variable domains of antibody heavy and light chains associated with each other) and scFv (comprising an antibody V_H domain linked to a V_L domain by a flexible peptide linker) fragments. Where the antigen binding site comprises a single variable domain, this may
5 be a heavy chain variable domain, most suitably a heavy chain variable domain derived from an immunoglobulin naturally devoid of light chains.

'Restricted conformational flexibility' relates to restriction of
10 movement of the antigen binding units about the backbone of the intervening polypeptide linker group.

The present invention may be more fully understood with reference to the following description, when read together with the accompanying
15 drawings. For convenience, an antigen binding protein comprising two single binding units is hereinafter referred to as a 'bi-head'.

BRIEF DESCRIPTION OF THE DRAWINGS

20 Figure 1 shows a nucleotide sequence of the PstI-BstEII insert of plasmid ~~pUR4640~~, encoding the heavy chain variable domain of an anti-RR6 antibody (denoted R9) from a llama.

Figure 2 shows the nucleotide sequence of the PstI-BstEII insert of
25 plasmid pUR4601, encoding the heavy chain variable domain of an anti-hCG antibody (denoted H14) from a llama.

Figure 3 shows a map of plasmid pUR4619.

30 Figure 4 shows the nucleotide sequence within plasmid pUR4619 which encodes an anti-hCG-anti-RR6 bispecific biheaded antigen-binding protein (denoted HI4-R9), missing the first 4 and last 3 amino acids.

35 Figure 5 shows the A405 signals of an ELISA to determine bispecificity of various HI4-R9 biheads.

Figure 6 shows the scores achieved in a rapid assay technology (RAT) format assay following the detection of 1 IU/ml hCG (human

chorionic gonadotrophin protein) with various anti-hCG-anti-RR6 bihead antigen binding proteins derived from a llama wherein the anti-hCG and anti-RR6 fragments are linked as follows (see Example 1.5):

5

Number	Linker	
1	no linker (directly attached)	
2	G-T-S-G-S	(SEQ. ID NO. 1)
3	S-S-S-A-S-A-S-S-A	(SEQ. ID NO. 2)
4	G-S-P-G-S-P-G	(SEQ. ID NO. 3)
5	A-T-T-T-G-S-S-P-G-P-T	(SEQ. ID NO. 4)
6	A-N-H-S-G-N-A-S	(SEQ. ID NO. 5)

Figure 7 shows a comparison of the sensitivity of detection of hCG in a RAT assay using various biheads (see Example 1.5).

10 DETAILED DESCRIPTION OF THE INVENTION

The invention is based on the unexpected finding that by using a polypeptide linking group conferring restricted conformational flexibility to link together antigen binding units, multivalent antigen binding proteins having advantageous binding affinity, as demonstrated by their increased sensitivity of diagnosis and detection, are obtained. Furthermore, constructs according to the invention may conveniently be produced at high yields economically and efficiently on a scale appropriate for industrial use.

20

As is apparent from the discussion of the background to the invention above, to the extent that multivalent antigen binding constructs comprising separate binding units linked together have been described at all in the literature, the linking means has been provided by flexible peptide groups. Flexibility of conformation in the linker group has been considered desirable in order to allow the multivalent construct to assume the correct orientation to allow simultaneous binding of multiple antigens.

30 Surprisingly, the present inventors have found that by restricting the conformational flexibility of the linking polypeptide group,

multivalent antigen binding constructs having improved binding affinity may be obtained. This is entirely contrary to the teaching in the art that the linking group should desirably be flexible.

- 5 The invention is applicable to the preparation of multivalent antigen binding constructs comprising antigen binding units where the antigen binding site is formed through association of antibody light and heavy chain variable domains. Preferably, however, the constructs prepared according to the invention comprise a plurality of single domain
- 10 binding units, more particularly a plurality of heavy chain variable domains derived from an immunoglobulin naturally devoid of light chains such as may be obtained from lymphoid cells, especially peripheral blood lymphocytes, bone marrow cells or spleen cells derived from Camelids as described in WO 94/04678 (Casterman et al)
- 15 discussed above. An advantage of using single domain binding units which are heavy chain variable domains derived from Camelids is that they can readily and conveniently be produced economically on a large scale, for example, using a transformed lower eukaryotic host, as described in WO 94/25591 (Unilever), described above.

20

- ~~Bivalent forms, that is having two antigen binding sites, of the~~
multivalent antigen binding proteins prepared according to the invention are preferred but it will be appreciated that higher multivalent forms, which are also encompassed in the present
- 25 invention, may find application under suitable circumstances, for example where more than two antigens are required to bind, for example in processes for scavenging molecules from solution or processes where close proximity of molecules form the basis of an assay.

- 30 Structural features which may suitably be incorporated into the linking polypeptide group in order to achieve the effect of restricting conformational flexibility according to the purposes of the invention would readily suggest themselves to those skilled in the art.

35

Accordingly, in one embodiment, the linker group preferably comprises one or more proline residues.

Without wishing to be bound by theory, it is generally thought that the presence of a proline residue in a peptide sequence encourages the amino acid backbone of the peptide to adopt a beta-turn structural configuration, with the peptide backbone changing direction about the proline residue. Linker groups comprising other sequence features which promote the formation of a beta-turn configuration in the peptide backbone, such as peptide linkers containing valine residues or constrained residues such as 8-bicyclic and 5,9-bicyclic tripeptide units (see, for example, Johannesson et al, J. Med. Chem., 42, 601-608 (1999)), may also suitably find application in the present invention.

In another embodiment, peptide linker groups derived from naturally occurring proteins such as cell wall proteins (CWP), in particular, CWP1, or cellobishydrolases (CBH), such as CBH1P, which serve to restrict conformational flexibility or linker groups showing at least 50% homology thereto as determined by the ALIGN program of Dayhoff et al (1983), Methods Enzymol., 91, 524-545, may also suitably be used according to the invention.

Peptide linker groups which encode a glycosylation binding site and/or ~~are resistant to proteolytic attack may also suitably be employed.~~ Here, the presence of a carbohydrate attached to the amino acid residues has the effect of restricting the flexibility of the peptide backbone.

Conveniently, the polypeptide linking group according to the invention comprises from 4 to 30 amino acid residues, preferably from 5 to 15 amino acid residues.

Preferred polypeptide linking groups according to the invention comprise an amino acid sequence selected from:

S-S-S-A-S-A-S-S-A, (SEQ. ID NO. 2)

G-S-P-G-S-P-G, (SEQ. ID NO. 3)

A-T-T-T-G-S-S-P-G-P-T (SEQ. ID NO. 4)

It will be appreciated that although the invention has been described primarily by reference to antigen binding proteins, it is equally applicable to proteins comprising other binding units as described

above. References to antigen binding proteins will accordingly be understood to refer also to such other proteins unless the context dictates otherwise.

- 5 Multivalent antigen binding proteins according to the invention may be prepared by transforming a host by incorporating a gene encoding the polypeptide as set forth above and expressing said gene in said host.

10 Suitably the host or hosts may be selected from prokaryotic bacteria, such as Gram-negative bacteria, for example *E. coli*, and Gram-positive bacteria, for example *B. subtilis* or lactic acid bacteria, lower eukaryotes such as yeasts, for example belonging to the genera *Saccharomyces*, *Kluyveromyces*, *Hansenula* or *Pichia*, or moulds such as those belonging to the genera *Aspergillus* or *Trichoderma*.

15 Preferred hosts for use in connection with the present invention are the lower eukaryotic moulds and yeasts.

20 Techniques for synthesising genes, incorporating them into hosts and expressing genes in hosts are well known in the art and the skilled person would readily be able to put the invention into effect using common general knowledge.

25 Methods for producing antibody fragments or functionalised fragments thereof derived from the heavy chain immunoglobulin of *Camelidae* using a transformed lower eukaryotic host are described, for example in patent application WO 94/25591 and such techniques may suitably be applied to prepare constructs according to the present invention.

30 Proteins according to the invention may be recovered and purified using conventional techniques such as affinity chromatography, ion exchange chromatography or gel filtration chromatography.

35 The activity of the multivalent binding proteins according to the invention may conveniently be measured by standard techniques known in the art such as enzyme-linked immunoadsorbant assay (ELISA), radioimmune assay (RIA) or by using biosensors.

The following examples are produced by way of illustration only.

Techniques used for the manipulation and analysis of nucleic acid materials were performed as described in Sambrook et al, Molecular Cloning, Cold Spring Harbor Press, New York, 2nd Ed., (1989) unless otherwise indicated.

5

Restriction sites are underlined.

HC-V denotes heavy chain variable domain.

EXAMPLES

10

Example 1 Self Assembling Llama Bi-heads Containing Linker Peptides on Latex to Assay hCG

1.1 Construction of Llama Bi-heads with Various Linkers

15

a) Induction of humeral immune responses in llama

20

Male llamas were immunised with a water in oil emulsion (1:9 V/V, antigen in water: Specol (Bokhout et al, Vet. Immunol. Immunopath., 2: 491-500 (1981)) subcutaneously and intramuscularly. Per immunisation site 0.75-1.5 ml water in oil emulsion was inoculated containing 100:µg antigen. The antigens used were: hCG (Sigma), azo-dye RR6 (ICI) which was coupled to BSA via its reactive triazine group. Immunisations were performed according to the following time table: The second immunisation was performed three weeks after the first. The third was performed two weeks after the second immunisation. The immune response was followed by antigen specific ELISAs.

25

30 The anti-RR-6 response was measured by using Nunc Covalink plates, which were coated with the azo-dye. After incubation with (diluted) serum samples, the bound llama antibodies were detected via a incubation with poly-clonal rabbit-anti-llama antiserum (obtained via immunising rabbits with llama immunoglobulins which were purified via
35 ProtA and ProtG columns; ID-DLO), followed by an incubation with swine-anti-rabbit immunoglobulins (Dako) conjugated with alkaline phosphatase. Finally the alkaline phosphatase enzyme-activity was determined after incubation with p-nitro-phenyl phosphate and the optical density was measured at 405nm. The anti-hCG response, was

measured in essentially the same way using Nunc maxi-sorb plates coated with hCG.

b) Cloning, expressing and screening of llama HC-V fragments

5

i) Isolation of gene fragments encoding llama HC-V domains

From an immunised llama a blood sample of about 200ml was taken and an enriched lymphocyte population was obtained via Ficoll (Pharmacia) discontinuous gradient centrifugation. From these cells, total RNA was isolated by acid guanidium thiocyanate extraction (e.g. via the method described by Chomczynski and Sacchi, Analytical Biochemistry, 162: 156-159 (1987). After first strand cDNA synthesis (e.g. with the Amersham first strand cDNA kit), DNA fragments encoding HC-V fragments and part of the long or short hinge region were amplified by PCR using specific primers:

20 V_H - 2B 5'-AGGTSMARCTGCAGSAGTCWGG-3' (SEQ. ID NO. 6)
 S = C and G, M = A and C, R = A and G, W = A and T,

25 Lam-07 5'-AACAGTTAAGCTTCCGCTTGCGGCCGCGGAGCTGGGGTCTTCGCTGTGGTGCG-3'
 (short hinge) (SEQ. ID NO. 7)

30 Lam-08 5'-AACAGTTAAGCTTCCGCTTGCGGCCGCTGGTTGTGGTTTTGGTGTCTTGGGTT-3'
 (long hinge) (SEQ. ID NO. 8)

Upon digestion of the PCR fragments with *Pst*I (coinciding with codon 4 and 5 of the HC-V domain, encoding the amino acids L-Q) and *Bst*EII (located at the 3'-end of the HC-V gene fragments; coinciding with the amino acid sequence Q-V-T), the DNA fragments with a length between 300 and 400bp (encoding the HC-V domain, but lacking the first three and the last three codons) were purified via gel electrophoresis and isolation from the agarose gel.

40 ii) Construction of *Saccharomyces cerevisiae* expression plasmids encoding llama HC-V domains

Plasmids pUR4547 and pUR4548 are *Saccharomyces cerevisiae* episomal expression plasmids, derived from pSY1 (Harmsen et al., Gene, 125: 115-123, (1993). From pSY1 the *Pst*I site, located in front of the GAL7 promoter was removed after partial digestion with *Pst*I, incubation with Klenow fragment and subsequent blunt end ligation. After transformation the desired plasmid could be selected on the basis of restriction pattern analysis. Subsequently, the *Bst*EII site in the *Leu2* selection marker was removed by replacing the about 410bp *Afl*III/*Pfl*MI fragment with a corresponding fragment in which the *Bst*EII site was removed via a three step PCR mutagenesis, using the primers:

PCR-A:

15 BOLI 1 5'-GGGAATTCCAATAGGTGGTTAGCAATCG (SEQ. ID NO. 9)
(*Pfl*MI)
BOLI 4 5'-GACCAACGTGGTCGCCTGGCAAACG (SEQ. ID NO. 10)
(*Bst*EII)

PCR-B:

20 BOLI 3 5'-CGTTTTGCCAGGCGACCACGTTGGTC (SEQ. ID NO. 11)
(*Bst*EII)

25 BOLI 2 5'-CCCCAAGCTTACATGGTCTTAAGTTGGCGT (SEQ. ID NO. 12)
(*Afl*III)

30 PCR-A was performed with primers BOLI 1 and BOLI 4 and resulted in an about 130bp fragment with the *Pfl*MI restriction site at the 3'-end and the inactivated *Bst*EII site at the 5'-end. PCR-B was performed with primers BOLI 2 and BOLI 3 and resulted in an about 290bp fragment with the *Afl*III site at the 5'-end. The third PCR was with the fragments obtained from reaction A and B, together with the primers BOLI 1 and BOLI 2.

35 Finally, the about 1.8kb *Sac*I-*Hind*III fragment was replaced with synthetic fragments, having sequences as presented below, resulting the plasmids pUR4547 and pUR4548, respectively.

- 12 -

- *SacI*/*HindIII* fragment of pUR4547

SacI (SEQ. ID NO. 13)
 GAGCTCATCACACAAACAAACAAACAAATGATGCTTTTGCAAGCCTTCCCTT
 5 1 -----+-----+-----+-----+----- 54
 CTCGAGTAGTGTGTTTGTGTTTGTGTTTACTACGAAAACGTTCCGAAGGGAA
 M M L L Q A F L F
 |→ SUC2 ss (SEQ. ID NO. 14)

10 *PstI*
 TTCCTTTTGGCTGGTTTTGCAGCCAAATATCTGCGCAGGTGCAGCTGCAGG
 55-----+-----+-----+-----+----- 105
 AAGGAAAACCGACCAAACGTCGGTTTTATAGACGCGTCCACGTCCGACGTCC
 L L A G F A A K I S A Q V Q L Q E
 15 |→

BstEII *HindIII*
 AGTCATAATGAGGGACCCAGGTACCGTCTCCTCATAATGACTTAAGCTT
 106-----+-----+-----+-----+----- 155
 20 TCAGTATTACTCCCTGGGTCCAGTGGCAGAGGAGTATTACTGAATTTCGAA
 E S * * G T Q V T V S S * *
 HC-V cassette ←|

and

25

- *SacI*/*HindIII* fragment of pUR4548

SacI (SEQ. ID NO. 15)
 GAGCTCATCACACAAACAAACAAACAAATGATGCTTTTGCAAGCCTTCCCTT
 30 1 -----+-----+-----+-----+----- 54
 CTCGAGTAGTGTGTTTGTGTTTGTGTTTACTACGAAAACGTTCCGAAGGAAA
 M M L L Q A F L F
 |→ SUC2 ss (SEQ. ID NO. 16)

35 *PstI*
 TCCTTTTGGCTGGTTTTGCAGCCAAATATCTGCGCAGGTGCAGCTGCAGG
 55-----+-----+-----+-----+----- 105
 AGGAAAACCGACCAAACGTCGGTTTTATAGACGCGTCCACGTCCGACGTCC
 L L A G F A A K I S A Q V Q L Q E
 40 |→

BstEII
 AGTCATAATGAGGGACCCAGGTACCGTCTCCTCAGAACAAAACTCATC
 106-----+-----+-----+-----+----- 155
 45 TCAGTATTACTCCCTGGGTCCAGTGGCAGAGGAGTCTTGTTTTTGAGTAG
 S * * G T Q V T V S S E Q K L I
 HC-V cassette ←|→ myc tail

- 13 -

HindIII

TCAGAAGAGGATCTGAATTAATGACTTAAGCTT
 156-----+-----+-----+----- 188
 AGTCTTCTCCTAGACTTAATTACTGAATTCGAA
 5 S E E D L N * *
 ←|

Both plasmids contain the GAL7 promoter and PGK terminator sequences as well as the invertase (SUC2) signal sequence. In both plasmids the

10 DNA sequence encoding the SUC2 signal sequence is followed by the first 5 codons, (encoding Q-V-Q-L-Q) of the HC-V domain (including the *Bst*II site), a stuffer sequence, the last six codons (encoding Q-V-T-V-S-S) of the HC-V domain. In pUR4547, this is followed by two stop codons, an *Afl*III and *Hind*III site. In pUR4548, this sequence is
 15 followed by eleven codons encoding the myc-tag, two stop codons, an *Afl*III and *Hind*III site.

Plasmids pUR4547 and pUR4548 were deposited under the Budapest Treaty at the Centraal Bureau voor Schimmelcultures, Baarn on 18th August
 20 1997 with deposition numbers: CBS 100012 and CBS 100013, respectively. In accordance with Rule 28(4) EPC, or a similar arrangement from a state not being a contracting state of the EPC, it is hereby requested that a sample of such deposit, when requested, will be submitted to an expert only.

25 Upon digesting pUR4548 with *Pst*I and *Bst*EII, the about 6.4kb vector fragment was isolated and ligated with the *Pst*I-*Bst*EII fragments of about 350bp obtained as described above. After transformation of *S. cerevisiae*, via electroporation, transformants were selected from
 30 minimal medium agar plates (comprising 0.7% yeast nitrogen base, 2% glucose and 2% agar, supplemented with the essential amino acids and bases).

iii) Screening for antigen specific HC-V domains
 35

For the production of llama HC-V fragments with myc-tail, individual transformants were grown overnight in selective minimal medium (comprising 0.7% yeast nitrogen base, 2% glucose, supplemented with the essential amino acids and bases) and subsequently diluted ten
 40 times in YPGal medium (comprising 1% yeast extract, 2% bacto pepton

and 5% galactose). After 24 and 48 hours of growth, the culture supernatant of the colonies was analysed by ELISA for the presence of HC-V fragments which specifically bind to the antigens hCG, RR6 in essential the same way as described above. In this case, however, the presence of specifically bound HC-V fragments was detected by incubation with monoclonal anti-myc antibodies, followed by incubation with poly-clonal rabbit-anti-mouse conjugate with alkaline phosphatase. In this way a number of anti-hCG and anti-RR6 HC-V fragments were isolated, which are:

10

anti-RR6:

R9 pUR4640 (see Figure 1) (SEQ. ID NO. 17, 18)

15 anti-hCG (alpha unit):

H14 pUR4601 (see Figure 2) (SEQ. ID NO. 19, 20)

20

c) Production of llama HC-V biheads by *S. cerevisiae*

~~1) Construction of episomal expression plasmids encoding anti-~~
hCG/anti-RR6 bispecific biheads

25 In the anti-hCG HC-V fragment H14 (anti-alpha-subunit), the *Pst*I site was removed and a *Xho*I site was introduced via PCR, using the primers:

MPG158WB

5'-GAATTAAGCGGCCGCCAGGTGAAACTGCTCGAGTCWGGGGGA-3' (SEQ. ID NO. 21)

30

and

MPG159WB

35 3'-CCCTGGGTCCAGTGGCAGAGGAGTGGCAGAGGAGTCTTGTTT-5' (SEQ. ID NO. 22)

In this way the sequence:

(SEQ. ID NO. 23)

XhoI

CAG	GTG	AAA	CTG	<u>CTC</u>	<u>GAG</u>	TCW	GGG
O	V	K	L	L	E	S	G

(SEQ. ID NO. 24)

Upon digesting the PCR fragments with *XhoI* and *BstEII*, the about 330bp fragments were purified via agarose gel electrophoresis and isolation from the gel. The fragments were cloned into pUR4421 (see Example 1 in WO 94/25591) which was digested with the same enzymes, resulting in pJS2 (H14). Subsequently, the about 420bp *EagI* -*HindIII* fragment of pJS2 was isolated and ligated in the about 6.6kb *EagI*- *HindIII* vector fragment of the pSY1 plasmid of which the *PstI* and *BstEII* sites were removed as described above. The resulting plasmid pJS7, was digested with *BstEII* and *HindIII*, after which the purified vector fragment was religated in the presence of a synthetic linker having the following sequence:

```

BstEII                               PstI                               HindIII
<-                               MPG 160 WB (49)                               -> (SEQ ID NO. 25)
GGTCACCGTCTCCTCACAGGTGCAGCTGCAGGAGTCACTGTAATGACTTAAGCTT
-----+-----+-----+-----+-----+-----+-----+----- 55
CCAGTGGCAGAGGAGTGTCCACGTCGACGTCCTCAGTGACATTACTGAATTTCGAA
      <-                               MPG 161 WB (48)                               ->
V  T  V  S  S  Q  V  Q  L  Q  E  S  L  *  *  L  K  L
                                           (SEQ. ID NO. 26)

```

resulting in plasmid pJS9. Finally, the plasmid was digested with *Pst*I and *Hind*III, after which the purified vector fragments of about 7.0kb were ligated with the *Pst*I -*Hind*III fragments of about 350bp of pUR4638 and pUR4640, encoding an anti-RR6 HC-V fragment denoted R9 followed by the myc-tail. The resulting *S. cerevisiae* episomal expression plasmid pUR4619 encodes a anti-hCG-anti-RR6 bispecific bihead preceded by the SUC2 signal sequence and followed by the myc-tail.

pUR4619: SUC2 - H14 - R9 - myc (see Figures 3-4/SEQ. ID NO. 27, 28)

Upon digesting these plasmids with *Xho*I and partially with *Bst*EII, *Xho*I-*Bst*EII fragments of about 0.7kb can be isolated and subsequently
5 cloned into the vector fragment of pUR4547 (digested with the same enzymes). In this way biheads can be obtained without the myc tail.

It will be appreciated that expression vectors can be constructed in which different promoter systems, e.g. the constitutive GAPDH promoter
10 or different signal sequences, e.g. the mating factor prepro sequence are used.

ii) Production of the HC-V biheads

15 After introducing the expression plasmid pUR4619 into *S. cerevisiae* via electroporation, transformants were selected from minimal medium agar plates as described in part b(ii) above. For the production of biheads, the transformants were grown overnight in selective minimal medium and subsequently diluted ten times in YPGal medium. After 24
20 and 48 hours of growth, samples were taken for Western blot analysis. ~~For the immuno-detection of the produced biheads via Western blot~~ analysis, monoclonal anti-myc antibodies were used, followed by incubation with poly-clonal rabbit-anti-mouse conjugate with alkaline phosphatase.

25

d) Anti-hCG/anti-RR6 bispecific biheads containing a linker peptide

i) Construction of *S. cerevisiae* episomal expression plasmids encoding anti-hCG/anti-RR6 bispecific biheads containing a
30 linker peptide

Between the H14 and the R9 encoding DNA fragments synthetic linkers were introduced encoding different linker peptides. To this end the about 50 bp long *Bst*EII-*Hind*III fragment of pJS7 (see Example 1 c(i)
35 above) was replaced by an about 50 bp long *Bst*EII-*Hind*III fragment having the following sequence:

MVaJA

BstEII *XbaI* *DraIII* *PstI* *HindIII*
 5' **GTCACCGTCTCTAGATGGCCACCAGGTGCAGCTGCAGGAGTCAACTTA** 3' (SEQ. ID NO. 29)

5 MVbJA

3' **GCAGAGATCTACCGGTGGTCCACGTCGAGCTCCTCAGTTGAATTCCA** 5'
 (SEQ. ID NO. 30)

10 This resulted in pSJ7a. In this plasmid the about 20 bp *PstI-HindIII* fragment was replaced with the about 370 bp *PstI-HindIII* fragment encoding the anti-RR6 HC-V fragment R9 and/or with the myc-tail of pUR4640 (see Example 1 c(i)) and resulting in pSJ7b.

15 Upon digesting plasmid pSJ7b with *XbaI* and *DraIII* the about 7 kb vector fragment was ligated with five synthetic oligo nucleotide linker fragments presented below:

MV01JA 5' CTAGTGGTACTTCCGGTTCCCAG 3' (SEQ. ID NO. 31)

20 MV02JA 3' ACCATGAAGGCCAAGG 5' (SEQ. ID NO. 32)

S G T S G S Q

25 MV03JA 5' CTAGTTCTTCATCTGCTTCTGCCTCTTCAGCCCAG 3' (SEQ. ID NO. 33)

MV04JA 3' AAGAAGTAGACGAAGACGGAGAAGTCGG 5' (SEQ. ID NO. 34)

S S S S A S A S S A Q

30 MV05JA 5' CTAGTGGTTCTCCAGGTTCCACCAGGTCAG 3' (SEQ. ID NO. 35)

MV06JA 3' ACCAAGAGGTCCAAGTGGTCCA 5' (SEQ. ID NO. 36)

S G S P G S P G Q

35

- 18 -

MV07JA 5' CTAGTGCTACTACAACCTGGTTCTTCACCAGGTCCAACCTCAG 3' (SEQ. ID NO. 37)

5 MV08JA 3' ACGATGATGTTGACCAAGAAGTGGTCCAGGTTGA 5' (SEQ. ID NO. 38)

S A T T T G S S P G P T Q

MV09JA 5' CTAGTGCTAATCATTCTGGTAATGCTTCTCAG 3' (SEQ. ID NO. 39)

10 MV10JA 3' ACGATTAGTAAGACCATTACGAAGA 5' (SEQ. ID NO. 40)

S A N H S G N A S Q

15 The oligonucleotide linker fragments encode the last amino acid of the N-terminal HC-V fragment (S) and the first amino acid of the C-terminal HC-V fragment, intersected by the connecting linker peptide. This resulted in plasmids pUR5330 to 5334, respectively.

20 After transformation of *S. cerevisiae* with these plasmids, the production levels of the biheads were determined via Western blot analysis and a anti-hCG ELISA using anti-myc mAb for detection of the bound bihead (see Example 1 b(iii)). Production levels are presented in Table 2 below:

25 Table 2

Plasmid	Linker	Production level (mg/l)
pUR4619	None	11
pUR5330	S-G-T-S-G-S-Q	36
pUR5331	S-S-S-S-A-S-A-S-S-A-Q	49
pUR5332	S-G-S-P-G-S-P-G-Q	33
pUR5333	S-A-T-T-T-G-S-S-P-G-P-T-Q	56
pUR5334	S-A-N-H-S-G-N-A-S-Q	51

30 The production levels of the biheads in which the two HC-V domains are separated by a linker peptide (consisting of between 5 and 11 amino acids) were found to be 3 to 5 times higher as found for the bihead in which the two HC-V fragments are connected without a peptide linker.

Finally, the bispecificity of the biheads was demonstrated as follows:

PINs coated with hCG were incubated with (diluted) medium samples.

- 5 Subsequently, the PINs were incubated with a RR6-alkaline phosphatase conjugate, in which the azo-dye RR6 was coupled to the alkaline phosphatase via its reactive triazine group. Finally, the alkaline phosphatase enzyme activity was determined after incubation of the PINs with p-nitro-phenyl phosphate and the optical density was
10 measured at 405nm (see Figure 5).

1.2 Purification of Llama Bi-heads with Various Linkers from *S. cerevisiae* Culture Media

- 15 A 5 ml column of recombinant Protein A Fast Flow Sepharose (Amersham Pharmacia Biotech) was equilibrated by washing with 10 column volumes of wash buffer (10 mM potassium phosphate, pH 6), at a flow rate of 2 ml/min. The bi-head fermentation broth was loaded at 2 ml/min in an upwards direction. After loading, the column was washed with wash
20 buffer until the OD₂₈₀ reached the baseline. Elution was carried out with a linear gradient of 0 - 40 mM citric acid pH 2.5 in the reverse direction, collecting 4 ml fractions into tubes containing 400 µl of a neutralising agent (1M Tris.Cl, pH 8.5) in order to minimise the effects of the acid. Peak fractions were checked for purity by
25 running on a 12% SDS-PAGE Ready Gel (Bio-Rad) under standard denaturing conditions. Staining was with GelCode Blue (Pierce & Warriner). The fractions were concentrated using Macrosep centrifugal concentrators (3 kDa molecular weight cut-off, Pall Filtron Corp.) then buffer exchanged into 10mM potassium phosphate, pH 6 using PD-10
30 columns (Amersham Pharmacia Biotech). The final purity of the sample was determined by carrying out a UV scan from 400 - 220 nm and using the value at 280 nm to determine an accurate concentration. The samples were then aliquoted into vials, frozen, freeze dried and stored until required.

35

1.3 Preparation of a Reactive Red 6-Bovine Serum Albumin Conjugate

A solution of Reactive Red 6 (RR6) was made up at 10 mg/ml in phosphate buffered saline (PBS). A solution of bovine serum albumin

(BSA) was made up at 10 mg/ml in PBS. 200 µl of the RR6 solution was added to 800 µl of the BSA solution and the resulting solution was mixed in an end over end rotary mixer for 2 hours at room temperature. RR6 that had conjugated to BSA was separated from free RR6 by addition of the reaction mixture (1 ml) to a PD10 column (Pharmacia) previously washed with 10 ml of PBS containing 0.1% sodium azide (PBSA). The column was then eluted by addition of PBSA (5 ml) and 1 ml aliquots were collected. The RR6-BSA conjugate eluted in fractions 4 and 5. These were pooled and the concentration of protein was determined using a BCA protein test and the concentration adjusted to 2 mg/ml with PBSA.

1.4 Adsorption of Latex with Reactive Red 6-Bovine Serum Albumin Conjugate

15

Duke blue latex was adsorbed with the RR6-BSA conjugate as follows:

To 950 µl of 10 mM borate buffer, 0.01 % merthiolate, pH 8.5 (buffer B) a 50 µl aliquot of Duke blue latex (10 % solids) was added and mixed by inverting. The diluted latex was centrifuged at 8000 g for 10 minutes at room temperature, the supernatant removed and the pellet vortexed briefly. The pellet was resuspended in 900 µl of buffer B and to this 100 µl of the previously prepared RR6-BSA conjugate was added. The latex solution was sonicated for 10 s using a sonic probe. The solution containing the latex was mixed for 2 h at room temperature and then centrifuged (8000 g, 10 min at room temperature). The latex pellet was washed by resuspending in 1 ml of buffer B and centrifuged once more (8000 g, 10 min at room temperature). The pellet was then resuspended in 1 ml buffer B ready for use.

30

1.5 Analysis of Llama Bi-head Self Assembling on Reactive Red 6-Bovine Serum Albumin Adsorbed Latex

The llama bi-heads were tested by self assembling onto RR6-BSA adsorbed latex and detection of hCG in a rapid assay technology (RAT) format. This was performed by mixing the llama bi-head (5 µl of a 0.1 mg/ml solution) with RR6-BSA adsorbed latex (5 µl of 0.1 % solids) in 10 µl of PBSA to which hCG (5 µl of various concentrations) was added.

The resulting solution was added to the bottom of a nitrocellulose strip (6 mm wide x 30 mm long) on which a monoclonal antibody recognising hCG had been adsorbed by plotting in a line (2.5 mg/ml) mid way up the strip. The latex-bi-head-hCG solution was allowed to
5 flow up the nitrocellulose strip by capillary action and the strip was then washed by applying PBSA (25 µl) to the bottom of the strip. The amount of latex, captured at the plotted antibody line on the nitrocellulose strip, was quantified by measuring the absorbance through the strip.

10

Figure 6 shows that the llama bi-heads with linkers 3, 4 and 5 gave the highest response in RAT assays. These linkers are structurally more ordered than the comparative examples, flexible linkers 2 and 6 and result in more hCG and more latex captured in the assay. The more
15 ordered linkers promote the correct orientation of the binding domains to achieve more optimal binding than when no linker is used. Linker 3 is derived from CWP1 and Linker 5 from CBH1P.

20

Synthetic linkers with some order (linker 4 containing 2 proline residues) can offer increased sensitivity in assays than those with little order (linker 2). Figure 7 shows that the bi-head with linker
4 can detect lower amounts (50 mIU/ml) of hCG than the bi-head with linker 2 and, hence, give a more sensitive assay for hCG.

25

SEQUENCE LISTING

5 <110> UNILEVER PLC
UNILEVER N.V

<120> ANTIGEN BINDING PROTEINS

10 <130> T3077

<140>
<141>

15 <160> 40

<170> PatentIn Ver. 2.1

20 <210> 1
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
25 <223> Description of Artificial Sequence:LINKER

<400> 1
Gly Thr Ser Gly Ser
1 5

30

<210> 2
<211> 9
<212> PRT
35 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LINKER

40 <400> 2
Ser Ser Ser Ala Ser Ala Ser Ser Ala
1 5

45 <210> 3
<211> 7
<212> PRT
<213> Artificial Sequence

50 <220>
<223> Description of Artificial Sequence:LINKER

<400> 3
Gly Ser Pro Gly Ser Pro Gly
55 1 5

<210> 4

- 23 -

<211> 11
<212> PRT
<213> Artificial Sequence

5 <220>
<223> Description of Artificial Sequence:PRIMER

<400> 4
Ala Thr Thr Thr Gly Ser Ser Pro Gly Pro Thr
10 1 5 10

15 <210> 5
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PRIMER

20 <400> 5
Ala Asn His Ser Gly Asn Ala Ser
1 5

25 <210> 6

30 <211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PRIMER

35 <400> 6
aggtsmarct gcagsagtcw gg 22

40 <210> 7
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PRIMER

45 <400> 7
aacagttaag cttccgcttg cggccgcgga gctggggtct tcgctgtggt gcg 53

50 <210> 8
<211> 53
<212> DNA
<213> Artificial Sequence

55 <220>
<223> Description of Artificial Sequence:PRIMER

<400> 8

- 24 -

aacagttaag cttccgcttg cggccgctgg ttgtggtttt ggtgtcttgg gtt 53

5 <210> 9
<211> 28
<212> DNA
<213> Artificial Sequence

10 <220>
<223> Description of Artificial Sequence:PRIMER

<400> 9
gggaattcca ataggtggtt agcaatcg 28

15

<210> 10
<211> 26
<212> DNA
<213> Artificial Sequence
20 <220>
<223> Description of Artificial Sequence:PRIMER

25 <400> 10
gaccaacgtg gtcgcctggc aaaacg 26

30 <210> 11
<211> 26
<212> DNA
<213> Artificial Sequence

35 <220>
<223> Description of Artificial Sequence:PRIMER

<400> 11
cgttttgccca ggcgaccacg ttggtc 26

40 <210> 12
<211> 30
<212> DNA
<213> Artificial Sequence

45 <220>
<223> Description of Artificial Sequence:PRIMER

50 <400> 12
ccccaagctt acatggtctt aagttggcgt 30

55 <210> 13
<211> 155
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PLASMID

- 25 -

CONSTRUCT

<220>

<221> CDS

5 <222> (3)..(140)

<400> 13

ga gct cat cac aca aac aaa caa aac aaa atg atg ctt ttg caa gcc 47
 Ala His His Thr Asn Lys Gln Asn Lys Met Met Leu Leu Gln Ala
 10 1 5 10 15

ttc ctt ttc ctt ttg gct ggt ttt gca gcc aaa ata tct gcg cag gtg 95
 Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys Ile Ser Ala Gln Val
 20 25 30

15

cag ctg cag gag tca taa tga ggg acc cag gtc acc gtc tcc tca 140
 Gln Leu Gln Glu Ser Gly Thr Gln Val Thr Val Ser Ser
 35 40 45

20 taatgactta agctt 155

<210> 14

<211> 46

25 <212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence:PLASMID
 CONSTRUCT

30 <400> 14

Ala His His Thr Asn Lys Gln Asn Lys Met Met Leu Leu Gln Ala Phe
 1 5 10 15

35 Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys Ile Ser Ala Gln Val Gln
 20 25 30

Leu Gln Glu Ser Gly Thr Gln Val Thr Val Ser Ser
 35 40 45

40

<210> 15

<211> 188

<212> DNA

<213> Artificial Sequence

45

<220>

<223> Description of Artificial Sequence:PLASMID
 CONSTRUCT

50

<220>

<221> CDS

<222> (3)..(173)

<400> 15

55 ga gct cat cac aca aac aaa caa aac aaa atg atg ctt ttg caa gcc 47
 Ala His His Thr Asn Lys Gln Asn Lys Met Met Leu Leu Gln Ala
 1 5 10 15

- 26 -

```

      ttc ctt ttc ctt ttg gct ggt ttt gca gcc aaa ata tct gcg cag gtg   95
      Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys Ile Ser Ala Gln Val
                20                      25                      30

5    cag ctg cag gag tca taa tga ggg acc cag gtc acc gtc tcc tca gaa   143
      Gln Leu Gln Glu Ser          Gly Thr Gln Val Thr Val Ser Ser Glu
                35                      40                      45

      caa aaa ctc atc tca gaa gag gat ctg aat taatgactta agctt         188
10   Gln Lys Leu Ile Ser Glu Glu Asp Leu Asn
                50                      55

```

```

      <210> 16
15   <211> 57
      <212> PRT
      <213> Artificial Sequence
      <223> Description of Artificial Sequence:PLASMID
            CONSTRUCT

20   <400> 16
      Ala His His Thr Asn Lys Gln Asn Lys Met Met Leu Leu Gln Ala Phe
          1              5              10              15

25   Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys Ile Ser Ala Gln Val Gln
            20              25              30

      Leu Gln Glu Ser          Gly Thr Gln Val Thr Val Ser Ser Glu Gln
            35              40              45

30   Lys Leu Ile Ser Glu Glu Asp Leu Asn
            50              55

```

```

35   <210> 17
      <211> 342
      <212> DNA
      <213> Artificial' Sequence

40   <220>
      <223> Description of Artificial Sequence:PLASMID
            CONSTRUCT

      <220>
45   <221> CDS
      <222> (1)..(342)

      <400> 17
50   cag gtg cag ctg cag gag tca ggg gga ggc ttg gtg cag gct ggg gag   48
      Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Glu
          1              5              10              15

      tct ctg aaa ctc tcc tgt gca gcc tct gga aac acc ttc agt ggc ggc   96
55   Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Asn Thr Phe Ser Gly Gly
          20              25              30

      ttc atg ggc tgg tac cgc cag gct cca ggg aag cag cgc gag ttg gtc   144
      Phe Met Gly Trp Tyr Arg Gln Ala Pro Gly Lys Gln Arg Glu Leu Val

```

- 27 -

```

          35          40          45
gca acc att aat agt aga ggt atc aca aac tat gca gac ttc gtg aag 192
Ala Thr Ile Asn Ser Arg Gly Ile Thr Asn Tyr Ala Asp Phe Val Lys
5          50          55          60

ggc cga ttc acc atc tcc aga gac aat gcc aag aag aca gtg tat ttg 240
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Lys Thr Val Tyr Leu
10         65          70          75          80

gaa atg aac agc ctg gaa cct gaa gac acg gcc gtt tat tac tgt tac 288
Glu Met Asn Ser Leu Glu Pro Glu Asp Thr Ala Val Tyr Tyr Cys Tyr
          85          90          95

15 act cac tac ttc aga tcc tac tgg ggt cag ggg acc cag gtc acc gtc 336
Thr His Tyr Phe Arg Ser Tyr Trp Gly Gln Gly Thr Gln Val Thr Val
          100          105          110

20 tcc tca 342
Ser Ser

<210> 18
<211> 114
25 <212> PRT
<213> Artificial Sequence
<223> Description of Artificial Sequence: PLASMID
CONSTRUCT

30 <400> 18
Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Glu
1          5          10          15

35 Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Asn Thr Phe Ser Gly Gly
20          25          30

Phe Met Gly Trp Tyr Arg Gln Ala Pro Gly Lys Gln Arg Glu Leu Val
35          40          45

40 Ala Thr Ile Asn Ser Arg Gly Ile Thr Asn Tyr Ala Asp Phe Val Lys
50          55          60

Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Lys Thr Val Tyr Leu
65          70          75          80

45 Glu Met Asn Ser Leu Glu Pro Glu Asp Thr Ala Val Tyr Tyr Cys Tyr
85          90          95

50 Thr His Tyr Phe Arg Ser Tyr Trp Gly Gln Gly Thr Gln Val Thr Val
100          105          110

Ser Ser

55 <210> 19
<211> 351
<212> DNA
<213> Artificial Sequence

```

- 28 -

<220>

<223> Description of Artificial Sequence: PLASMID
CONSTRUCT

5

<220>

<221> CDS

<222> (1)..(351)

10

<400> 19

cag gtg cag ctg cag gag tca ggg gga gga ttg gtg cag gcg ggg ggc 48

Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly

1 5 10 15

15

tct ctg aga ctc tcc tgt gca gcc tct gga cgc acc ggc agt acg tat 96

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Thr Gly Ser Thr Tyr

20 25 30

20

gac atg ggc tgg ttc cgc cag gct cca ggg aag gag cgt gag tct gta 144

Asp Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val

35 40 45

25

gca gct att aac tgg gat agt gcg cgc aca tac tat gca agc tcc gtg 192

Ala Ala Ile Asn Trp Asp Ser Ala Arg Thr Tyr Tyr Ala Ser Ser Val

50 55 60

30

agg ggc cga ttc acc atc tcc aga gac aac gcc aag aag acg gtg tat 240

Arg Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Lys Thr Val Tyr

65 70 75 80

ctg caa atg aac age etg aaa eet gag gac acg gcc gtt tat acc tgt 288

Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Thr Cys

85 90 95

35

ggc gcg ggg gaa ggt ggt act tgg gac tcc tgg ggc cag ggg acc cag 336

Gly Ala Gly Glu Gly Gly Thr Trp Asp Ser Trp Gly Gln Gly Thr Gln

100 105 110

40

gtc acc gtc tcc tca 351

Val Thr Val Ser Ser

115

<210> 20

45

<211> 117

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence: PLASMID
CONSTRUCT

50

<400> 20

Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly

1 5 10 15

55

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Thr Gly Ser Thr Tyr

20 25 30

Asp Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val

- 29 -

35 40 45

Ala Ala Ile Asn Trp Asp Ser Ala Arg Thr Tyr Tyr Ala Ser Ser Val
50 55 60

5 Arg Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Lys Thr Val Tyr
65 70 75 80

10 Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Thr Cys
85 90 95

Gly Ala Gly Glu Gly Gly Thr Trp Asp Ser Trp Gly Gln Gly Thr Gln
100 105 110

15 Val Thr Val Ser Ser
115

20 <210> 21
<211> 43
<212> DNA
<213> Artificial Sequence

25 <220>
<223> Description of Artificial Sequence:PRIMER

<400> 21
gaattaagcg gccgcccagg tgaaactgct cgagtcwggg gga 43

30

<210> 22
<211> 42
<212> DNA
<213> Artificial Sequence

35 <220>
<223> Description of Artificial Sequence:PRIMER

<400> 22
40 ccctgggtcc agtggcagag gagtggcaga ggagtcttgt tt 42

45 <210> 23
<211> 24
<212> DNA
<213> Artificial Sequence

50 <220>
<223> Description of Artificial Sequence:PRIMER

<400> 23
caggtccagc tgcaggagtc tggg 24

55 <210> 24
<211> 24
<212> DNA
<213> Artificial Sequence

- 30 -

<220>

<223> Description of Artificial Sequence:PRIMER

5 <400> 24
caggtgaaac tgctcgagtc wggg 24

10 <210> 25

<211> 55

<212> DNA

<213> Artificial Sequence

15 <220>

<223> Description of Artificial Sequence:LINKER;DOUBLE
STRANDED

20 <220>

<221> CDS

<222> (2)..(55)

25 <400> 25
g gtc acc gtc tcc tca cag gtg cag ctg cag gag tca ctg taa tga ctt 49
Val Thr Val Ser Ser Gln Val Gln Leu Gln Glu Ser Leu Leu
1 5 10 15

30 aag ctt 55
Lys Leu

35 <210> 26

<211> 18

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence:LINKER;DOUBLE
STRANDED

40 <400> 26
Val Thr Val Ser Ser Gln Val Gln Leu Gln Glu Ser Leu Leu
1 5 10 15

Lys Leu

45

<210> 27

<211> 672

<212> DNA

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence:PLASMID
CONSTRUCT

55

<220>

<221> CDS

<222> (1)..(672)

<400> 27

	ctc	gag	tca	ggg	gga	gga	ttg	gtg	cag	gcg	ggg	ggc	tct	ctg	aga	ctc	48
	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Ala	Gly	Gly	Ser	Leu	Arg	Leu	
5	1				5					10					15		
	tcc	tgt	gca	gcc	tct	gga	cgc	acc	ggc	agt	acg	tat	gac	atg	ggc	tgg	96
	Ser	Cys	Ala	Ala	Ser	Gly	Arg	Thr	Gly	Ser	Thr	Tyr	Asp	Met	Gly	Trp	
				20					25					30			
10	ttc	cgc	cag	gct	cca	ggg	aag	gag	cgt	gag	tct	gta	gca	gct	att	aac	144
	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val	Ala	Ala	Ile	Asn	
				35				40					45				
15	tgg	gat	agt	gcg	cgc	aca	tac	tat	gca	agc	tcc	gtg	agg	ggc	cga	ttc	192
	Trp	Asp	Ser	Ala	Arg	Thr	Tyr	Tyr	Ala	Ser	Ser	Val	Arg	Gly	Arg	Phe	
		50					55					60					
20	acc	atc	tcc	aga	gac	aac	gcc	aag	aag	acg	gtg	tat	ctg	caa	atg	aac	240
	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Lys	Thr	Val	Tyr	Leu	Gln	Met	Asn	
	65					70				75					80		
25	agc	ctg	aaa	cct	gag	gac	acg	gcc	gtt	tat	acc	tgt	ggc	gcg	ggg	gaa	288
	Ser	Leu	Lys	Pro	Glu	Asp	Thr	Ala	Val	Tyr	Thr	Cys	Gly	Ala	Gly	Glu	
				85					90					95			
	ggt	ggt	act	tgg	gac	tcc	tgg	ggc	cag	ggg	acc	cag	gtc	acc	gtc	tcc	336
	Gly	Gly	Thr	Trp	Asp	Ser	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	
				100				105					110				
30	tca	cag	gtg	cag	ctg	cag	gag	tca	ggg	gga	ggc	ttg	gtg	cag	gct	ggg	384
	Ser	Gln	Val	Gln	Leu	Gln	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Ala	Gly	
		115					120					125					
35	gag	tct	ctg	aaa	ctc	tcc	tgt	gca	gcc	tct	gga	aac	acc	ttc	agt	ggc	432
	Glu	Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Asn	Thr	Phe	Ser	Gly	
		130					135					140					
40	ggc	ttc	atg	ggc	tgg	tac	cgc	cag	gct	cca	ggg	aag	cag	cgc	gag	ttg	480
	Gly	Phe	Met	Gly	Trp	Tyr	Arg	Gln	Ala	Pro	Gly	Lys	Gln	Arg	Glu	Leu	
	145				150				155						160		
45	gtc	gca	acc	att	aat	agt	aga	ggt	atc	aca	aac	tat	gca	gac	ttc	gtg	528
	Val	Ala	Thr	Ile	Asn	Ser	Arg	Gly	Ile	Thr	Asn	Tyr	Ala	Asp	Phe	Val	
				165				170						175			
	aag	ggc	cga	ttc	acc	atc	tcc	aga	gac	aat	gcc	aag	aag	aca	gtg	tat	576
	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Lys	Thr	Val	Tyr	
				180				185					190				
50	ttg	gaa	atg	aac	agc	ctg	gaa	cct	gaa	gac	acg	gcc	gtt	tat	tac	tgt	624
	Leu	Glu	Met	Asn	Ser	Leu	Glu	Pro	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
		195					200					205					
55	tac	act	cac	tac	ttc	aga	tcc	tac	tgg	ggt	cag	ggg	acc	cag	gtc	acc	672
	Tyr	Thr	His	Tyr	Phe	Arg	Ser	Tyr	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	
		210					215					220					

- 32 -

<210> 28
<211> 224
<212> PRT
<213> Artificial Sequence
5 <223> Description of Artificial Sequence: PLASMID
CONSTRUCT

<400> 28
10 Leu Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly Ser Leu Arg Leu
1 5 10 15

Ser Cys Ala Ala Ser Gly Arg Thr Gly Ser Thr Tyr Asp Met Gly Trp
20 25 30

15 Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val Ala Ala Ile Asn
35 40 45

Trp Asp Ser Ala Arg Thr Tyr Tyr Ala Ser Ser Val Arg Gly Arg Phe
50 55 60

20 Thr Ile Ser Arg Asp Asn Ala Lys Lys Thr Val Tyr Leu Gln Met Asn
65 70 75 80

Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Thr Cys Gly Ala Gly Glu
25 85 90 95

Gly Gly Thr Trp Asp Ser Trp Gly Gln Gly Thr Gln Val Thr Val Ser
100 105 110

30 Ser Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly
115 120 125

Glu Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Asn Thr Phe Ser Gly
130 135 140

35 Gly Phe Met Gly Trp Tyr Arg Gln Ala Pro Gly Lys Gln Arg Glu Leu
145 150 155 160

Val Ala Thr Ile Asn Ser Arg Gly Ile Thr Asn Tyr Ala Asp Phe Val
40 165 170 175

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Lys Thr Val Tyr
180 185 190

45 Leu Glu Met Asn Ser Leu Glu Pro Glu Asp Thr Ala Val Tyr Tyr Cys
195 200 205

Tyr Thr His Tyr Phe Arg Ser Tyr Trp Gly Gln Gly Thr Gln Val Thr
50 210 215 220

<210> 29
<211> 48
<212> DNA
55 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: SYNTHETIC

INSERT

5 <400> 29
gtcaccgtct ctagatggcc accaggtgca gctgcaggag tcaactta 48

10 <210> 30
<211> 47
<212> DNA
<213> Artificial Sequence

15 <220>
<223> Description of Artificial Sequence:SYNTHETIC
INSERT

20 <400> 30
gcagagatct accggtgggc cacgtcgagc tcctcagttg aattcga 47

20 <210> 31
<211> 23
<212> DNA
<213> Artificial Sequence

25 <220>
~~<223> Description of Artificial Sequence:LINKER~~

30 <400> 31
ctagtgttac ttccggttcc cag 23

35 <210> 32
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LINKER

40 <400> 32
accatgaagg ccaagg 16

45 <210> 33
<211> 35
<212> DNA
<213> Artificial Sequence

50 <220>
<223> Description of Artificial Sequence:LINKER

55 <400> 33
ctagtcttc atctgcttct gcctcttcag cccag 35

<210> 34
<211> 28
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:LINKER

5 <400> 34
aagaagtaga cgaagacgga gaagtcgg 28

10 <210> 35
<211> 29
<212> DNA
<213> Artificial Sequence

15 <220>

<223> Description of Artificial Sequence:LINKER

20 <400> 35
ctagtgggtc tccaggttca ccaggtcag 29

<210> 36
<211> 22
<212> DNA

25 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:LINKER

30 <400> 36
~~accaagaggt ccaagtggc ca~~ 22

35 <210> 37
<211> 41
<212> DNA
<213> Artificial Sequence

40 <220>

<223> Description of Artificial Sequence:LINKER

<400> 37
ctagtgtac tacaactggt tcttcaccag gtccaactca g 41

45 <210> 38
<211> 34
<212> DNA
<213> Artificial Sequence

50 <220>

<223> Description of Artificial Sequence:LINKER

<400> 38

55 acgatgatgt tgaccaagaa gtgggccagg ttga 34

<210> 39

<211> 32
<212> DNA
<213> Artificial Sequence

5 <220>
<223> Description of Artificial Sequence:LINKER

<400> 39
ctagtgtctaa tcattctggt aatgcttctc ag 32
10

<210> 40
<211> 25
<212> DNA
15 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LINKER
20 <400> 40
acgattagta agaccattac gaaga 25



CLAIMS

1. Use of a polypeptide group, the amino acid sequence of which group confers restricted conformational flexibility, as a
5 linking group to link binding units in a multivalent binding protein.

2. ~~Use according to claim 1 wherein the polypeptide linking group comprises from 4 to 30 amino acid residues.~~

10

3. Use according to claim 1 or 2 wherein the linking group comprises one or more proline residues.

4. Use according to claim 1 or 2 wherein the linking group
15 comprises an amino acid sequence selected from:

S-S-S-A-S-A-S-S-A,

G-S-P-G-S-P-G, or

A-T-T-T-G-S-S-P-G-P-T.

20

5. A multivalent binding protein comprising a plurality of binding units linked by means of intervening polypeptide linker groups, the amino acid sequence of which linker group confers restricted conformational flexibility.

25

6. A protein according to claim 5 wherein the binding units comprise heavy chain variable domains derived from an immunoglobulin naturally devoid of light chains.

30 7. A protein according to claim 5 or claim 6 wherein the antigen binding units comprise heavy chain variable domains derived from a Camelid immunoglobulin.

8. A protein according to any one of claims 5 to 7 comprising a
35 bivalent antigen binding protein.

9. A protein according to any one of claims 5 to 8 wherein the linker group comprises from 4 to 30 amino acid residues.

10. A protein according to any one of claims 5 to 9 wherein the linker group comprises one or more proline residues.
11. A protein according to any one of claims 5 to 9 wherein the linker group comprises an amino acid sequence selected from:
- 5 S-S-S-A-S-A-S-S-A,
 G-S-P-G-S-P-G, or
 A-T-T-T-G-S-S-P-G-P-T.
- 10
12. Nucleotide sequences encoding for a multivalent binding protein of any one of claims 5 to 11.
13. An expression vector comprising a nucleotide sequence according to claim 12.
- 15
14. A host cell transformed with a vector according to claim 13.

ABSTRACT

Use of a polypeptide linker group, the amino acid sequence of which group confers restricted conformational flexibility, as a linking
5 group to link binding units, preferably antigen binding units comprising heavy chain variable domains derived from an immunoglobulin naturally devoid of light chains, in a multivalent binding protein.



PstI

CAGGTGCAGCTGCAGGAGTCAGGGGGAGGCTTGGTGCAGGCTGGGGAGTCTCTGAAACTCTCCTGTGCAGCCTCTGGAAACACCTTCAGT
1 -----+-----+-----+-----+-----+-----+-----+-----+-----+ 90
GTCCACGTCGACGTCCTCAGTCCCCCTCCGAACCACGTCCGACCCCTCAGAGACTTTGAGAGGACACGTCCGAGACCTTTGTGGAAGTCA
Q V Q L Q E S G G G L V Q A G E S L K L S C A A S G N T F S
[-> CDR I

KpnI

GGCGGCTTCATGGGCTGGTACCGCCAGGCTCCAGGGAAGCAGCGCGAGTTGGTCGCAACCATTAAATAGTAGAGGTATCACAACTATGCA
91 -----+-----+-----+-----+-----+-----+-----+-----+ 180
CCGCGGAAGTACCCGACCATGGCGGTCCGAGGTCCCTTCGTCGCGCTCAACCAGCGTTGGTAATTATCATCTCCATAGTGTGTTGATACGT
G G F M G W Y R Q A P G K Q R E L V A T I N S R G I T N Y A
[-] [-> CDR II

EagI

GACTTCGTGAAGGGCCGATTCAACATCTCCAGAGACAATGCCAAGAAGACAGTGTATTTGGAAATGAACAGCCTGGAACCTGAAGACACG
181 -----+-----+-----+-----+-----+-----+-----+-----+ 270
CTGAAGCACTTCCCGGCTAAGTGGTAGAGGTCTCTGTTACGGTTCTTCTGTGCACATAAACCTTTACTTGTGCGGACCTTGGACTTCTGTGC
D F V K G R F T I S R D N A K K T V Y L E M N S L E P E D T
[-]

BstEII

GCCGTTTATTACTGTTACACTCACTACTTCAGATCCTACTGGGGTCAGGGGACCCAGGTCACCGTCTCCTCA
271 -----+-----+-----+-----+-----+-----+-----+-----+ 342
CGGCAAATAATGACAATGTGAGTGATGAAGTCTAGGATGACCCAGTCCCCTGGGTCCAGTGGCAGAGGAGT
A V Y Y C Y T H Y F R S Y W G Q G T Q V T V S S
[-> CDR III [-]

Figure 1.

PstI

CAGGTGCAGCTGCAGGAGTCAGGGGGAGGATTGGTGCAGGCGGGGGGCTCTCTGAGACTCTCCTGTGCAGCCTCTGGACGCACCGGCAGT

1 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 90

GTCCACGTGACGTCTCAGTCCCCCTCCTAACCACGTCCGCCCCCGAGAGACTCTGAGAGGACACGTCCGAGACCTGCGTGGCCGTCA

Q V Q L Q E S G G G L V Q A G G S L R L S C A A S G R T G S

ACGTATGACATGGGCTGGTTCGGGCGAGGCTGCACCGAAGCAGCCTCAGTCTGTAGCAGCTATTAAGTGGGATAGTGGCGGCACATACTAT

91 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 180

TGCATACTGTACCCGACCAAGGCGGTCCGAGGTCCCTTCCTCGCACTCAGACATCGTCGATAATTGACCCCTATCACGCGCGTGTATGATA

T Y D M G W F R Q A P G K E R E S V A A I N W D S A R T Y Y

1-> CDR I <-1 1-> CDR II

EagI

GCAAGCTCCGTGAGGGGCGGATTACCATCTCCAGAGACAACGCCAAGAAGACGGTGTATCTGCAAATGAACAGCCTGAAACCTGAGGAC

181 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 270

CGTTCGAGGCACTCCCCGGCTAAGTGGTAGAGGTCTCTGTTGCGGTTCTTCTGCCACATAGACGTTTACTTGTGCGGACTTTGGACTCCTG

A S S V R G R F T I S R D N A K K T V Y L Q M N S L K P E D

<-1

BstEII

ACGGCCGTTTATACCTGTGGCGCGGGGAAGGTGGTACTTGGGACTCCTGGGGCCAGGGGACCCAGGTACCGTCTCCTCA

271 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 351

TGCGGGCAATATGGACACCGCGCCCCCTTCCACCATGAACCTGAGGACCCCGGTCCCCTGGGTCCAGTGGCAGAGGAGT

T A V Y T C G A G E G G T W D S W G Q G T Q V T V S S

1-> CDR III <-1

Figure 2.




Figure 3.

XhoI

CTCGAGTCAGGGGGAGGATTGGTGCAGGCGGGGGCTCTCTGAGACTCTCTGTGCAGCCTCTGGACGCACCGGCAGTACGTATGACATG
 1 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 90
 GAGCTCAGTCCCCCTCCTAACCACGTCCGCCCCCGAGAGACTCTGAGAGGACACGTCGGAGACCTGCGTGGCCGTCATGCATACTGTAC
 L E S G G G L V Q A G G S L R L S C A A S G R T G S T Y D M
 1-> CDR I

GGCTGGTTCGCCAGGCTCCAGGGAAGGAGCGTGAGTCTGTAGCAGCTATTAAGTGGGATAGTGC GCGCACATACTATGCAAGCTCCGTG
 91 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 180
 CCGACCAAGGCGGTCCGAGGTCCCTTCCTCGCACTCAGACATCGTCGATAATTGACCTATCACGCGCGTGTATGATACGTTTCGAGGCAC
 G W F R Q A P G K E R E S V A A I N W D S A R T Y Y A S S V
 <-1 1-> CDR II

EagI

AGGGGCGGATTACCATCTCCAGAGACAACGCCAAGAAGACGGTGTATCTGCAAATGAACAGCCTGAAACCTGAGGACACGGCCGTTTAT
 181 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 270
 TCCCCGGCTAAGTGGTAGAGGTCTCTGTGCGGTTCTTCTGCCACATAGACGTTTACTTGTGCGACTTTGGACTCCTGTGCCGCAATA
 R G R F T I S R D N A K K T V Y L Q M N S L K P E D T A V Y
 <-1

BstEII

↓

PstI

ACCTGTGGCGCGGGGAAGGTGGTACTTGGGACTCCTGGGGCCAGGGGACCCAGGTACCCGTCTCCTCACAGGTGCAGCTGCAGGAGTCA
 271 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 360
 TGGACACCGCGCCCCCTTCCACCATGAACCTGAGGACCCCGTCCCTGGGTCCAGTGGCAGAGGAGTGTCCACGTGACGCTCCTCAGT
 T C G A G E G G T W D S W G Q G T Q V T V S S Q V Q L Q E S
 1-> CDR III <-1

KpnI

GGGGGAGGCTTGGTGCAGGCTGGGGAGTCTCTGAAACTCTCCTGTGCAGCCTCTGGAACACCTTCAGTGGCGGCTTCATGGGCTGGTAC
 361 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 450
 CCCCCCTCCGAACCACGTCGACCCCTCAGAGACTTTGAGAGGACACGTCGGAGACCTTTGTGGAAGTCACCGCCGAAGTACCCGACCATG
 G G G L V Q A G E S L K L S C A A S G N T F S G G F M G W Y
 1-> CDR I <-1

CGCCAGGCTCCAGGGAAGCAGCGGAGTTGGTGCACCAACCTTAATAGTAGAGGTATCACAACTATGCAGACTTCGTGAAGGGCCGATTC
 451 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 540
 GCGGTCCGAGGTCCCTTCGTGCGCTCAACCAGCGTTGGTAATTATCATCTCCATAGTGTGATACGTCTGAAGCACTTCCCGGCTAAG
 R Q A P G K Q R E L V A T I N S R G I T N Y A D F V K G R F
 1-> CDR II <-1


EagI

ACCATCTCCAGAGACAATGCCAAGAAGACAGTGTATTTGGAAATGAACAGCCTGGAACCTGAAGACACGGCCGTTTATTACTGTTACACT
 541 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 630
 TGGTAGAGGTCTCTGTTACGGTTCTTCTGTACATAAACCTTTACTTGTGCGGACCTTGGACTTCTGTGCCGCAATAATGACAATGTGA
 T I S R D N A K K T V Y L E M N S L E P E D T A V Y Y C Y T

BstEII

CACTACTTCAGATCCTACTGGGGTCAGGGGACCCAGGTACCC
 631 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 672
 GTGATGAAGTCTAGGATGACCCAGTCCCTGGGTCCAGTGG
 H Y F R S Y W G Q G T Q V T
 1-> CDR III <-1

Figure 4.



pUR4619 pUR5330 pUR5331 pUR5332
pUR5333 pUR5334 blanc

Figure 5.




Figure 6.




Figure 7.



2
1
2

